

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number
WO 02/02731 A1

(51) International Patent Classification⁷: **C11D 17/00**,
3/00, 1/62, 3/02, 3/50, 3/20

(21) International Application Number: **PCT/US01/20353**

(22) International Filing Date: **27 June 2001 (27.06.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
09/607,033 **29 June 2000 (29.06.2000)** **US**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: **COLGATE-PALMOLIVE COMPANY**
[US/US]; 300 Park Avenue, New York, NY 10022 (US).

Published:
— with international search report

(72) Inventors: **GRANDMAIRE, Jean-Paul**; Rue Sous le Chateau 45, B-4821 Andrimont (BE). **HERMOSILLA, Anita**; Rue d'Heur 31, B-4340 Othee (BE).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(74) Agent: **LIEBERMAN, Bernard**; Colgate-Palmolive Company, 909 River Road, P.O. Box 1343, Piscataway, NJ 08855-1343 (US).



WO 02/02731 A1

(54) Title: **MULTI-PHASE CLEAR FABRIC SOFTENING COMPOSITION**

(57) Abstract: An aqueous multi-phase liquid fabric softener composition comprised of at least two phases, all of said phases being transparent or translucent, said fabric softener composition comprising: (a) from about 5% to about 45%, by weight, of a cationic softening material; (b) from about 5% to about 15%, by weight, of a solvent having a partition coefficient of log P of from about 0.70 to about 2.0; (c) from about 0.2% to about 1.5%, by weight, of a mineral electrolyte; (d) a water-immiscible oil-based perfume in an amount sufficient to provide the desired degree of fragrance; and (e) balance water whereby said multi-phase softener composition is capable of being converted to a milky macroemulsion upon dilution with water.

Multi-Phase Clear Fabric Softening Composition

This invention relates to rinse cycle fabric softener compositions. More particularly, it relates to an aqueous multi-phase liquid fabric softener composition comprising at least two transparent or translucent phases.

Multi-phase compositions which are transparent or clear are well-known in products intended for personal care such as bath oils and shower products, but not in fabric softening compositions. The various technologies which are used to obtain a clear multi-phase aesthetic effect generally rely on compositions containing oil and/or anionic surfactants that are generally incompatible with cationic surfactants and, particularly with the type of cationic surfactant required to provide a fabric softening composition which is both efficacious and transparent or clear.

Clear single phase fabric softening compositions are described in the patent literature. U.S. Patent Nos. 5,525,245 and 5,656,585 assigned to Colgate-Palmolive, and U.S. Patent No. 5,427,697 to Procter & Gamble describe single phase solutions or microemulsions which contain fabric softening ingredients. The microemulsions of the aforementioned '245 and '585 patents are said to be converted to macroemulsions having a turbid or milky appearance upon dilution with water in the rinse cycle thereby enabling fabric softening to occur.

While consumer preference often favors the aesthetic of clarity in fabric softener compositions, there remains as an object of the prior art the ability to provide a multi-phase fabric softener composition in which all phases are either clear or transparent or translucent, notwithstanding that such composition contains one or more cationic softening ingredients. In addition to the enhanced aesthetic properties, the resulting composition must also provide efficacious fabric softening when used as a rinse cycle softening composition.

Summary of the Invention

The objects cited above are satisfied by an aqueous multi-phase liquid fabric softener composition comprised of at least two phases, all of said phases being transparent or translucent, said fabric softener composition comprising:

- (a) from about 5% to about 45%, by weight, of a cationic softening material;
- (b) from about 5% to about 15%, by weight, of a solvent having a partition coefficient of log P of from about 0.70 to about 2.0;
- (c) from about 0.2% to about 1.5%, by weight, of a mineral electrolyte;


- (d) a water-immiscible oil-based perfume in a sufficient amount to provide the desired degree of fragrance; and
- (e) balance water whereby said multi-phase softener composition is capable of being converted to a milky macroemulsion upon dilution with water.

5 The present invention is predicated upon the discovery that the selection of a solvent and a mineral electrolyte in accordance with the invention when combined with a diester quaternary ammonium fabric softener as herein described provide an efficacious liquid fabric softener having at least two phases and wherein each phase manifests the desired aesthetic property of being transparent or translucent.

10 All of the ingredients of the softening composition must be normally liquid, namely, liquid at ambient room-temperature.

Detailed Description of the Invention

15 The cationic softening material for purposes of the invention is preferably a diester quaternary ammonium surfactant fabric softener selected from the group having the structural formulae as follows:



wherein each

20 A is independently C(O)O-R' or -O(O)-C-R';


R is a lower group having 1 to about 4 carbon atoms;

R' is an alkyl or alkenyl group having 8 to about 22 carbon atoms;


R'' is independently a lower alkyl radical having 1 to about 6 carbon atoms or hydroxyl alkyl group;

25 n is an integer having a value of 1 to about 3;

X⁻ is a softener compatible anion, preferably selected from the group consisting of a halide ion and methyl or ethyl sulfate; and



wherein B independently is A or (R)_n-A; and A, R, R'' and n are as defined above; and



wherein A, R, R'' and n are as defined above.

The preferred diester quaternary ammonium surfactant fabric softeners are represented by equation (I) above and are commercially available from Stepan Co. as Stepanex and from Kao Corp. as Tetranyl but can also be synthesized by the reaction of two moles of a fatty acid with a trialkanolamine optionally followed by alkoxylation and methylation with dimethyl sulfate or an alkyl halide such as, methyl iodide. In a preferred mode the fatty acid is oleic acid and ethylene oxide is used as the alkoxylation agent. For economical reasons it has been found that Soya fatty acids are a practical source for this purpose consisting of about 3% myristic acid, about 5% palmitic acid, about 5% palmitoleic acid, 1.5% stearic acid, 72.5% oleic acid and about 13% linoleic acid. Other sources of useful fatty acids are those obtained from the saponification of beef tallow, butter, corn oil, cottonseed oil, lard, olive oil, palm oil, peanut oil, cod liver oil, coconut oil and the like.

A preferred diester quaternary ammonium surfactant fabric softener is methyl bis[ethyl(oleyl)]-2-hydroxyethyl ammonium methyl sulfate. This quaternary ammonium compound is often referred to herein as dioleyl diesterquat (or "DODEQ").

Other diesters useful in the practice of this invention include:

methyl bis-[ethyl(coconut)]-2-hydroxyethyl ammonium methyl sulfate
 methyl bis-[ethyl(decyl)]-2-hydroxyethyl ammonium methyl sulfate
 methyl bis-[ethyl(dodecyl)]-2-hydroxyethyl ammonium methyl sulfate
 methyl bis-[ethyl(lauryl)]-2-hydroxyethyl ammonium methyl sulfate
 methyl bis-[ethyl(palmityl)]-2-hydroxyethyl ammonium methyl sulfate
 methyl bis-[ethyl(soft-tallow)]-2-hydroxyethyl ammonium methyl sulfate, and the like.

The designation of the terms coconut and beef tallow indicate mixtures of esters corresponding to the fatty acid source.

In the preparation of the diester quaternary ammonium surfactants, a certain amount of the triester homolog may be produced as an impurity. Unlike the diester, it is not soluble in water and has to be considered as an oil to be emulsified.

The term "oil-based perfume" is used herein in its ordinary sense to refer to and include any non water-soluble fragrant substance or mixture of substances including natural (i.e., obtained by extraction of flower, herb, blossom or plant), artificial (i.e., a mixture of

natural oils or oil constituents) and synthetic (i.e., a single or mixture of synthetically produced substance) odoriferous substances. Typically perfumes are complex mixtures of blends of various organic compounds, such as, esters, ketones, hydrocarbons, lactones, alcohols, aldehydes, ethers, aromatic compounds and varying amounts of essential oils (e.g.,
5 terpenes) such as from about 0% to about 80%, usually from about 10% to 70% by weight, the essential oils themselves being volatile odoriferous compounds and also serving to dissolve the other components of the perfume. The precise composition of the perfume has no particular effect on fabric softening so long as it meets the criteria of water immiscibility and pleasant odor. The level of perfume in the present softener compositions will generally
10 vary from about 0.1% to about 2.5%, by weight.

The solvents useful for the present invention are selected based on their octanol/water partition coefficient (P). This coefficient is the ratio between the equilibrium concentration of the particular solvent in octanol and in water. The coefficient is commonly provided in the technical literature as the logarithm (to the base 10) of P, namely, log P.

15 The value of log P of many solvents is often available from the solvent supplier or may be calculated, the "calculated log P" being referred to as C log P. For many solvents, the value of log P is reported and available from Daylight Chemical Information Systems Inc. (Daylight CIS), Irvine, California on the Pomona 92 database. Log P values are also conveniently calculated by the "C log P" program, also available from Daylight CIS. The C
20 log P values are determined based on the fragment approach of Hansch and Leo (cf., A. Leo, in Comprehensive Medicinal Chemistry, Vol. 4, C. Hansch, P.G. Sammens, J.B. Taylor and C.A. Ramsden Eds., p. 295, Pergamon Press, 1990, incorporated herein by reference).

Solvents generally useful for the present invention are those having a partition coefficient of log P of from about 0.70 to about 2.0. The particularly preferred range of log P
25 is from about 1.5 to about 2.0

Preferred solvents for use in the present invention are the following, the value of log P being indicated alongside in parentheses: dipropylene glycol n-propyl ether (0.994); tripropylene glycol n-butyl ether (1.896); dipropylene glycol n-butyl ether (1.523); tripropylene glycol n-butyl ether (1.9); and propylene glycol n-butyl (1.15). Other useful
30 solvents are diols and alcohols, such as, trimethyl pentane diol (1.24); pentanol (1.3); and hexanol (2.0).

Preferred mineral electrolytes suitable for use in this invention include: sodium sulfate; sodium nitrate; calcium chloride; ammonium sulfate; potassium chloride; potassium nitrate; calcium nitrate; magnesium nitrate; and magnesium sulfate.

An important aesthetic property of the present liquid fabric softening compositions is that following vigorous agitation of the composition, such as by hand shaking, the multi-phase product is temporarily converted to a milky macroemulsion. However, after a period of time ranging from about 5 minutes to about 12 hours, phase separation occurs although not as clear or transparent phases. After a period of about 5 hours to 72 hours, the original clear multiple phases reappear to restore the original aesthetic property of the composition. Also, when diluted in the rinse water, the multi-phase composition forms a macroemulsion thereby enabling the softening ingredients which are no longer solubilized in solution or microemulsion form to contact the fabrics in the rinse bath and provide the desired softening effect.

Example 1

A typical composition in accordance with the invention is comprised as follows:

Component	Wt. Percent
DODEQ ⁽¹⁾	21.8
Na ₂ SO ₄	0.5
Perfume	1.2
Pentanol	5.0
Water	71.4

⁽¹⁾ DODEQ refers to 90% Dioleyl Diesterquat plus 10% ethanol.

The above-described composition is characterized by two clear or transparent phases. Upon agitation the composition becomes turbid or milky. Following such agitation, phase separation occurs within about 3 minutes and the composition returns to its clear condition within about 5 hours.

Example 2

Compositions A-E were prepared as shown in Table 1 to compare the aesthetic appearance of compositions in accordance with the invention (B and C) with compositions outside the invention (A, D and E) all of the compositions being identical except for the selection of solvent. The value of log P for each solvent is indicated in parentheses.

Table 1					
Component	Composition A	Composition B	Composition C	Composition D	Composition E
Water	71.4 wt%	71.4 wt%	71.4 wt%	71.4 wt%	71.4 wt%
DODEQ	21.8	21.8	21.8	21.8	21.8
Perfume	1.2	1.2	1.2	1.2	1.2
Na ₂ SO ₄	0.25	0.25	0.25	0.25	0.25
Propanol (log P=0.3)	5	---	---	---	---
Pentanol (log P=1.3)	---	5	---	---	---
Hexanol (log P=2.0)	---	---	5	---	---
Octanol (log P=3.15)	---	---	---	5	---
Decanol (log P=3.32)	---	---	---	---	5
Appearance	Turbid	3 Clear Phases	3 Clear Phases	Turbid Gel	White Gel

As noted in Table 1, compositions B and C containing solvents having partition coefficients in accordance with the invention provided the desired aesthetic appearance of a clear multi-phase composition. In contrast thereto, compositions A, D and E containing solvents having values of log P outside the invention resulted in an undesired aesthetic appearance of turbidity associated with the formation of a macro-emulsion.

Example 3

Compositions F-H were prepared as shown in Table 2 to demonstrate the difference in aesthetic appearance provided by a composition in accordance with the invention (H) relative to compositions outside the invention (F and G) wherein the compositions differ only with respect to the selection of solvent.

Table 2			
Weight % of Components			
Component	Composition F	Composition G	Composition H
Water	71.4	71.4	71.4
DODEQ	21.8	21.8	21.8
Perfume	1.2	1.2	1.2
Na ₂ SO ₄	0.5	0.5	0.5
Propylene Glycol (log P = -1.4)	5	---	---
1-4 Butanediol (log P = -1.3)	---	5	---
Trimethyl Pentanediol (log P = 1.24)	---	---	5
Appearance	Turbid Gel	Turbid Gel	2 Clear Phases

As noted in Table 2, Composition H in accordance with the invention provided the desired aesthetic appearance in contrast to compositions F and G containing solvents not in accordance with the invention which were turbid or milky in appearance.

5

Example 4

Compositions I-M were prepared as shown in Table 3 using various propylene glycol ether solvents. The aesthetic appearance was noted for compositions in accordance with the invention (J, K, L and M) versus a composition outside the invention (I).

Table 3					
	Composition I	Composition J	Composition K	Composition L	Composition M
Water	7.14 wt%	7.14 wt%	7.14 wt%	7.14 wt%	7.14 wt%
DODEQ	21.8	21.8	21.8	21.8	21.8
Perfume	1.2	1.2	1.2	1.2	1.2
Na ₂ SO ₄	0.5	0.5	0.5	0.5	0.5
PnP (log P = 0.62) ¹	7.5	---	---	---	---
DPnP (log P = 0.99) ²	---	7.5	---	---	---
DPnB (log P = 1.52) ³	---	---	7.5	---	---
TPnB (log P = 1.9) ⁴	---	---	---	7.5	---
PnB (log P = 1.15) ⁵	---	---	---	---	7.5
Appearance	Turbid	2 Clear Phases	2 Clear Phases	2 Clear Phases	2 Clear Phases

10

¹ PnP refers to a Dowanol solvent which is propylene glycol n-propyl ether

² DPnP refers to a Dowanol solvent which is dipropylene glycol n-propyl ether

³ DPnB refers to a Dowanol solvent which is dipropylene glycol n-butyl ether

⁴ TPnB refers to tripropylene glycol n-butyl ether

15

⁵ PnB refers to propylene glycol n-butyl ether

As noted in Table 3, Composition I which is characterized by a solvent having a value of log P outside of the invention did not provide the desired aesthetic appearance of clarity and multiple phases. Compositions J, K, L and M manifested 2 clear phases.

20

Example 5

This Example compares the effect of using various electrolytes in two prototype compositions, I and II shown below:

25

Component	Composition I	Composition II
Water	71.4 wt%	71.4 wt%
DODEQ	16.67	16.67
Perfume	1.2	1.2
Electrolyte	0.5	0.9
TPnB	5	5

Mineral electrolytes (in accordance with the invention) and organic electrolytes (outside the invention) were used in prototype compositions I and II with the resulting aesthetic appearance being indicated as shown in Table 4 below:

5

Table 4					
Appearance of Compositions I and II with Various Electrolytes					
Mineral Electrolytes	Electrolyte	Composition I Examples 1-12		Composition II Examples 13-24	
	Na ₂ SO ₄	Ex. 1	2 Clear Phases	Ex. 13	2 Clear Phases
	NaCl	2	2 Clear Phases	14	2 Clear Phases
	NaNO ₃	3	2 Clear Phases	15	2 Clear Phases
	CaCl ₂	4	2 Clear Phases	16	2 Clear Phases
	MgCl ₂	5	2 Clear Phases	17	2 Clear Phases
	NH ₄ Sulfate	6	2 Clear Phases	18	2 Clear Phases
Organic Electrolytes					
	Quaternized TEA	7	Turbid	19	2 Clear Phases
	Na Benzoate	8	Turbid	20	2 Clear Phases
	Na Gluconate	9	Turbid	21	Turbid
	Na Citrate	10	Turbid	22	3 Clear Phases
	Na Tartrate	11	Turbid	23	Turbid
	Ca Formiate	12	Turbid	24	Turbid

As noted in Table 4, clear or transparent dual phase compositions were obtained with all of the mineral electrolytes tested in compositions I and II, all of which are compositions in accordance with the invention. The use of organic electrolytes (outside of the invention) resulted primarily in turbid or milky compositions, although in a few examples at the higher concentration level of electrolyte (Examples 19, 20 and 22), clear multiphase products were obtained.

15

Example 6

The softening performance of a composition in accordance with the invention was compared to that of a commercial rinse cycle softening composition. Specifically, composition L described in Example 4, Table 3 (containing 21.8% DODEQ, 90% of which is active softener ingredient) was evaluated for softening performance versus a commercial

concentrated fabric softening composition containing a comparable amount of softener ingredients (reference composition). The dosage for both compositions in the rinse cycle was 28 ml.

5 Test fabrics to be evaluated were first desized to remove the finish. The test fabrics were introduced into a standard fabric load and treated in a European washing machine under European detergent and water hardness conditions.

The washed fabrics were line dried and the test fabrics were then evaluated through pair testing using a panel of judges.

10 The resulting softness delivered by the composition of the invention (composition L) was judged to be essentially equivalent to the softness delivered by the reference composition.

Claims

What is claimed is:

- 5 1. An aqueous multi-phase liquid fabric softener composition comprised of at least two phases, all of said phases being transparent or translucent, said fabric softener composition comprising:
- 10 (a) from about 5% to about 45%, by weight, of a cationic softening material;
- (b) from about 5% to about 15%, by weight, of a solvent having a partition coefficient of log P of from about 0.70 to about 2.0;
- (c) from about 0.2% to about 1.5%, by weight, of a mineral electrolyte;
- (d) a water-immiscible oil-based perfume in an amount sufficient to provide the desired degree of fragrance; and
- 15 (d) balance water whereby said multi-phase softener composition is capable of being converted to a milky macroemulsion upon dilution with water.
2. An aqueous fabric softener composition in accordance with claim 1 wherein said solvent is an alcohol or a diol.
- 20 3. An aqueous fabric softener composition in accordance with claim 1 wherein said solvent is dipropylene glycol n-propyl ether.
4. An aqueous fabric softener composition in accordance with claim 1 wherein said solvent is dipropylene glycol n-butyl ether
- 25 5. An aqueous fabric softener composition in accordance with claim 1 wherein said solvent is tripropylene glycol n-butyl ether.
6. An aqueous fabric softener composition in accordance with claim 1 wherein said
- 30 solvent is a propylene glycol ether.
7. An aqueous fabric softener composition in accordance with claim 1 wherein the cationic softening material is a diester quaternary ammonium surfactant.

8. An aqueous fabric softener composition as in claim 7 wherein the diester is methyl bis[ethyl(oleyl)]-2-hydroxyethyl ammonium methyl sulfate.

5 9. An aqueous fabric softener composition as in claim 1 comprised of two transparent phases.

10. An aqueous phase fabric softener composition as in claim 1 comprised of three transparent phases.

10 11. A method of softening fabrics comprising the step of contacting said fabrics in a rinse bath with an effective amount of the fabric softener composition of claim 1.

INTERNATIONAL SEARCH REPORT

onal Application No
US 01/20353

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C11D17/00 C11D3/00 C11D1/62 C11D3/02 C11D3/50
C11D3/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 040 287 A (MOTYKA ET AL) 21 March 2000 (2000-03-21)	1,7-9,11
Y	column 2, line 18 -column 6, line 37 column 6, line 55 -column 7, line 1 claims	1-7,10
Y	US 5 656 585 A (GRANDMAIRE JEAN-PAUL ET AL) 12 August 1997 (1997-08-12) cited in the application column 1, line 65 -column 3, line 30 column 4, line 6 - line 25 claims	1-4,7
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

12 October 2001

Date of mailing of the international search report

18/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Serbetsoglou, A

INTERNATIONAL SEARCH REPORT

International Application No.

JS 01/20353

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 665 268 A (DEGUERTECHIN LOUIS OLDENHOVE ET AL) 9 September 1997 (1997-09-09)	1,2,5,10
A	column 6, line 40 - line 50 column 9, line 3 - line 33 claim 1	3,4,6
Y	US 5 958 863 A (KEYS ROBERT O) 28 September 1999 (1999-09-28) column 4, line 49 -column 11, line 20 column 12, line 33 - line 49 claims	1,6
A	WO 92 17523 A (PROCTER & GAMBLE) 15 October 1992 (1992-10-15) claims 1,6,7 page 14, line 9 -page 17, line 21 examples IV-XI page 28, line 29 - line 30	1,2,7-11

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No.
PCT/JP 01/20353

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6040287	A	21-03-2000	US 5856287 A	05-01-1999
			US 6040071 A	21-03-2000
			AU 696570 B2	10-09-1998
			AU 4577896 A	05-09-1996
			CA 2170699 A1	02-09-1996
			EP 0730023 A2	04-09-1996
			NZ 286025 A	24-04-1997
US 5656585	A	12-08-1997	US 5525245 A	11-06-1996
			AU 737508 B2	23-08-2001
			AU 3217297 A	07-01-1998
			BR 9709674 A	09-05-2000
			CA 2253509 A1	18-12-1997
			EP 0912709 A2	06-05-1999
			WO 9747723 A2	18-12-1997
			ZA 9704975 A	07-12-1998
			AU 691720 B2	21-05-1998
			AU 4525096 A	10-07-1996
			BR 9510472 A	26-05-1998
			CA 2208368 A1	27-06-1996
			CN 1173199 A	11-02-1998
			CZ 9701925 A3	12-11-1997
			EP 0801672 A1	22-10-1997
			FI 972647 A	18-08-1997
			HU 77479 A2	28-05-1998
			IL 116474 A	17-02-2000
			JP 10511146 T	27-10-1998
			NO 972874 A	13-08-1997
			NZ 300493 A	25-11-1998
			PL 321433 A1	08-12-1997
			RO 115174 B1	30-11-1999
			RU 2141998 C1	27-11-1999
			TR 960592 A2	21-07-1996
			WO 9619552 A1	27-06-1996
			ZA 9510746 A	18-06-1997
US 5665268	A	09-09-1997	NONE	
US 5958863	A	28-09-1999	US 5674832 A	07-10-1997
			AU 5671996 A	18-11-1996
			CA 2218725 A1	31-10-1996
			EP 0822859 A1	11-02-1998
			JP 11505555 T	21-05-1999
			WO 9633800 A1	31-10-1996
			US 5753079 A	19-05-1998
WO 9217523	A	15-10-1992	AU 1753292 A	02-11-1992
			CA 2106217 A1	29-09-1992
			CN 1066680 A	02-12-1992
			JP 6506251 T	14-07-1994
			MX 9201415 A1	01-10-1992
			NZ 242150 A	27-04-1995
			PT 100298 A	30-07-1993
			WO 9217523 A1	15-10-1992

THIS PAGE BLANK (USPTO)